

Example: Let  $X$  denote the number of girls born in the births of 4 babies, i.e. how many are girls??

Idea: Define  $X_j = 1$  if  $j$ th baby is a girl  
 $0$  otherwise

Notice: Always  $X = X_1 + X_2 + X_3 + X_4$ . It is just like counting on your fingers. E.g. if only babies 2 and 3 are girls, then

$$X_1 = 0, X_2 = 1, X_3 = 1, X_4 = 0 \quad X = 0 + 1 + 1 + 0 = 2$$

Notice  $E(X_j) = P(A_j)$  where  $A_j$  is the event that the  $j$ th baby is a girl.  
So  $E(X_j) = \frac{1}{2}$  for each  $j$ .

$$\text{So altogether } E(X) = E(X_1 + X_2 + X_3 + X_4) \\ = E(X_1) + E(X_2) + E(X_3) + E(X_4) = \frac{1}{2} + \frac{1}{2} + \frac{1}{2} + \frac{1}{2} = 2.$$

Same expected value we got as when we first treated the problem.

Also: notice we never computed the mass of  $X$ ,  
i.e. we never wrote down the numbers  $\frac{1}{16}, \frac{4}{16}, \frac{6}{16}, \frac{4}{16}, \frac{1}{16}$ .

So this method is easier, more straightforward, and more natural since we are used to counting occurrences (yes/no) on your fingers!