

Density and the CDF for Gamma random variables with parameters  $\lambda$  and  $r$   
The density for such a Gamma random variable is

$$f_X(x) = \frac{\lambda^r}{(r-1)!} x^{r-1} e^{-\lambda x},$$

for  $x > 0$ , and  $f_X(x) = 0$  otherwise.

Notice that in the  $r = 1$  case, we just have  $f_X(x) = \lambda e^{-\lambda x}$  just like it was for Exponential random variables, because a Gamma random variable with  $r = 1$  is exactly an Exponential random variable. So the  $r = 1$  case has to agree.

What about the CDF of a Gamma random variable? It is

$$F_X(x) = 1 - e^{-\lambda x} \sum_{j=0}^{r-1} \frac{(\lambda x)^j}{j!},$$

for  $x > 0$ , and  $F_X(x) = 0$  otherwise.

Again, let's check the  $r = 1$  case. If  $r = 1$ , then the CDF is  $F_X(x) = 1 - e^{-\lambda x}$ , and this agrees exactly with the CDF of an Exponential random variable, as it must, because again, a Gamma random variable with parameter  $r = 1$  is exactly an Exponential random variable.

Also the expected value of a Gamma random variable must be the sum of  $r$  copies the expected value of an Exponential random variable. In other words, if  $X$  is a Gamma random variable with parameters  $\lambda$  and  $r$ , then  $X$  has the same distribution as  $X_1 + X_2 + \dots + X_r$ , where the  $X_j$ 's are independent Exponential random variables. Therefore

$$E(X) = E(X_1 + X_2 + \dots + X_r) = E(X_1) + E(X_2) + \dots + E(X_r) = \frac{1}{\lambda} + \frac{1}{\lambda} + \dots + \frac{1}{\lambda} = \frac{r}{\lambda}.$$

Similarly, since the  $X_j$ 's are independent random variables (now we need the independence), we can also compute:

$$Var(X) = Var(X_1 + X_2 + \dots + X_r) = Var(X_1) + Var(X_2) + \dots + Var(X_r) = \frac{1}{\lambda^2} + \frac{1}{\lambda^2} + \dots + \frac{1}{\lambda^2} = \frac{r}{\lambda^2}.$$