

STAT/MA 41600
Practice Problems: October 3, 2014
Solutions by Mark Daniel Ward

1. Hungry customers.

a. The number X of people in the survey who are eating pizza is Hypergeometric with $M = 7$, $N = 12$, and $n = 3$. So the mass of X is $p_X(x) = \frac{\binom{7}{x}\binom{5}{3-x}}{\binom{12}{3}}$.

b. The values are

$$\begin{aligned}p_X(0) &= \frac{\binom{7}{0}\binom{5}{3-0}}{\binom{12}{3}} = 1/22, \\p_X(1) &= \frac{\binom{7}{1}\binom{5}{3-1}}{\binom{12}{3}} = 7/22, \\p_X(2) &= \frac{\binom{7}{2}\binom{5}{3-2}}{\binom{12}{3}} = 21/44, \\p_X(3) &= \frac{\binom{7}{3}\binom{5}{3-3}}{\binom{12}{3}} = 7/44.\end{aligned}$$

c. We can write the average as $(0)(1/22) + (1)(7/22) + (2)(21/44) + (3)(7/44) = 7/4$, but it is perhaps easier to note that each person in the survey eats pizza with probability $7/12$, so the average number of survey participants eating pizza is $7/12 + 7/12 + 7/12 = (3)(7/12) = 21/12 = 7/4$.

2. Harmonicas.

a. The number X of Deluxe harmonicas is Hypergeometric with $M = 7$, $N = 19$, and $n = 8$. So $\mathbb{E}(X) = (8)(7/19) = 56/19 = 2.9474$. This can also be seen by the fact that each harmonica selected has a probability $7/19$ of being a Deluxe harmonica.

b. Since the number X of Deluxe harmonicas is Hypergeometric with $M = 7$, $N = 19$, and $n = 8$, then $\text{Var}(X) = n\frac{M}{N}\left(1 - \frac{M}{N}\right)\left(\frac{N-n}{N-1}\right) = (8)(7/19)(1 - 7/19)(11/18) = 1232/1083 = 1.1376$.

c. Since the number X of Deluxe harmonicas is Hypergeometric with $M = 7$, $N = 19$, and $n = 8$, then $P(X = 5) = \frac{\binom{7}{5}\binom{12}{3}}{\binom{19}{8}} = 770/12597 = 0.0611$.

3. Granola bars.

a. The number X of chocolates I grab is Hypergeometric with $M = 6 + 8 = 14$, $N = 24$, and $n = 3$. So $P(X = 2) = \frac{\binom{14}{2}\binom{10}{1}}{\binom{24}{3}} = 455/1012 = 0.4496$.

b. The probability that strictly fewer than 2 are chocolate is $P(X = 0) + P(X = 1) = \frac{\binom{14}{0}\binom{10}{3}}{\binom{24}{3}} + \frac{\binom{14}{1}\binom{10}{2}}{\binom{24}{3}} = 15/253 + 315/1012 = 375/1012 = 0.3706$.

c. The average number of chocolate granola bars is $n(M/N) = 3(14/24) = 7/4$. This can also be determined by the fact that each bar is chocolate with probability $14/24$, so the average number of chocolates is $3(14/24) = 7/4$.

4. Superfans.

a. The exact probability is $\binom{60,000}{8} \left(\frac{1}{10,000}\right)^8 \left(\frac{9999}{10,000}\right)^{59992}$.

b. If $n = 60,000$ and $p = 1/10,000$, then the probability in part (a) is the probability that a Binomial n, p random variable is equal to 8. Now let X be Poisson with average $\lambda = np = 6$. Then $P(X = 8) = \frac{e^{-6}6^8}{8!}$.

c. We have $P(X = 8) = \frac{e^{-6}6^8}{8!} = 0.1033$.

5. Shoppers.

a. The exact probability is

$$= \binom{100,000}{0} \left(\frac{49,999}{50,000}\right)^{100,000} + \binom{100,000}{1} \left(\frac{1}{50,000}\right)^1 \left(\frac{49,999}{50,000}\right)^{99,999} \\ + \binom{100,000}{2} \left(\frac{1}{50,000}\right)^2 \left(\frac{49,999}{50,000}\right)^{99,998} + \binom{100,000}{3} \left(\frac{1}{50,000}\right)^3 \left(\frac{49,999}{50,000}\right)^{99,997}$$

b. If $n = 100,000$ and $p = 1/50,000$, then the probability in part (a) is the probability that a Binomial n, p random variable is 3 or less. Now let X be Poisson with average $\lambda = np = 2$. Then $P(X \leq 3) = \frac{e^{-2}2^0}{0!} + \frac{e^{-2}2^1}{1!} + \frac{e^{-2}2^2}{2!} + \frac{e^{-2}2^3}{3!}$.

c. We have $P(X \leq 3) = \frac{e^{-2}2^0}{0!} + \frac{e^{-2}2^1}{1!} + \frac{e^{-2}2^2}{2!} + \frac{e^{-2}2^3}{3!} = 0.8571$.